

# Effective Categoricity of Equivalence Structures

## FINAL DRAFT

**Wesley Calvert**

Department of Mathematics  
University of Notre Dame  
wcalvert@nd.edu

**Douglas Cenzer**

Department of Mathematics  
University of Florida  
cenzer@math.ufl.edu

**Valentina Harizanov**

Department of Mathematics  
George Washington University  
harizanv@gwu.edu

**Andrei Morozov**

Sobolev Institute of Mathematics  
morozov@math.nsc.ru

May 13, 2008

## 1 Introduction

We consider only countable structures for computable languages with universe  $\omega$ . We identify sentences with their Gödel codes. The *atomic diagram* of a structure  $\mathcal{A}$  for  $L$  is the set of all quantifier-free sentences in  $L_A$ ,  $L$  expanded by constants for the elements in  $A$ , which are true in  $\mathcal{A}$ . A structure is *computable* if its atomic diagram is computable. In other words, a structure  $\mathcal{A}$  is computable if there is an algorithm that determines for every quantifier-free formula  $\theta(x_0, \dots, x_{n-1})$  and every sequence  $(a_0, \dots, a_{n-1}) \in A^n$ , whether  $\mathcal{A} \models \theta(a_0, \dots, a_{n-1})$ . The *elementary diagram* of  $\mathcal{A}$  is the set of all sentences of  $L_A$  that are true in  $\mathcal{A}$ . A structure  $\mathcal{A}$  is *decidable* if its elementary diagram is computable. For  $n > 0$ , the *n-diagram* of  $\mathcal{A}$  is the set of all  $\Sigma_n$  sentences of  $L_A$  that are true in  $\mathcal{A}$ . A structure is *n-decidable* if its *n*-diagram is computable.

A computable structure  $\mathcal{A}$  is *computably categorical* if for every computable isomorphic copy  $\mathcal{B}$  of  $\mathcal{A}$ , there is a *computable* isomorphism from  $\mathcal{A}$  onto  $\mathcal{B}$ . For example, the ordered set of rational numbers is computably categorical, while

the ordered set of natural numbers is not. Moreover, Goncharov and Dzgoev [12] and Remmel [28] independently proved that a computable linear ordering is computably categorical if and only if it has only finitely many successors. They also established that a computable Boolean algebra is computably categorical if and only if it has finitely many atoms (see also LaRoche [21]). Miller [26] proved that no computable tree of height  $\omega$  is computably categorical. Lempp, McCoy, Miller and Solomon [22] characterized computable trees of finite height that are computably categorical. Nurtazin [27] and Metakides and Nerode [24] established that a computable algebraically closed field of finite transcendence degree over its prime field is computably categorical. Goncharov [9] and Smith [30] characterized computably categorical abelian  $p$ -groups as those that can be written in one of the following forms:  $(Z(p^\infty))^l \oplus G$  for  $l \in \omega \cup \{\infty\}$  and  $G$  finite, or  $(Z(p^\infty))^n \oplus G \oplus (Z(p^k))^\infty$ , where  $n, k \in \omega$  and  $G$  is finite. Goncharov, Lempp and Solomon [15] proved that a computable, ordered, abelian group is computably categorical if and only if it has finite rank. Similarly, they showed that a computable, ordered, Archimedean group is computably categorical if and only if it has finite rank.

For any computable ordinal  $\alpha$ , we say that a computable structure  $\mathcal{A}$  is  $\Delta_\alpha^0$  *categorical* if for every computable structure  $\mathcal{B}$  isomorphic to  $\mathcal{A}$ , there is a  $\Delta_\alpha^0$  isomorphism from  $\mathcal{A}$  onto  $\mathcal{B}$ . Lempp, McCoy, Miller and Solomon [22] proved that for every  $n \geq 1$ , there is a computable tree of finite height that is  $\Delta_{n+1}^0$ -categorical but not  $\Delta_n^0$ -categorical. We say that  $\mathcal{A}$  is *relatively computably categorical* if for every structure  $\mathcal{B}$  isomorphic to  $\mathcal{A}$ , there is an isomorphism that is computable relative to the atomic diagram of  $\mathcal{B}$ . Similarly, a computable  $\mathcal{A}$  is *relatively  $\Delta_\alpha^0$  categorical* if for every  $\mathcal{B}$  isomorphic to  $\mathcal{A}$ , there is an isomorphism that is  $\Delta_\alpha^0$  relative to the atomic diagram of  $\mathcal{B}$ . Clearly, a relatively  $\Delta_\alpha^0$  categorical structure is  $\Delta_\alpha^0$  categorical. We are especially interested in the case when  $\alpha = 2$ . McCoy [23] characterized, under certain restrictions, all  $\Delta_2^0$  categorical and relatively  $\Delta_2^0$  categorical linear orderings and Boolean algebras. For example, a computable Boolean algebra is relatively  $\Delta_2^0$  categorical if and only if it can be expressed as a finite direct sum  $c_1 \vee \dots \vee c_n$ , where each  $c_i$  is either atomless, an atom, or 1-atom. Using an enumeration result of Selivanov [29], Goncharov [10] showed that there is a computable structure that is computably categorical but not relatively computably categorical. Using a relativized version of Selivanov's enumeration result, Goncharov, Harizanov, Knight, McCoy, Miller and Solomon [13] showed that for each computable successor ordinal  $\alpha$ , there is a computable structure that is  $\Delta_\alpha^0$  categorical, but not relatively  $\Delta_\alpha^0$  categorical.

There are syntactical conditions that are equivalent to relative  $\Delta_\alpha^0$  categoricity. The conditions involve the existence of certain families of formulas, that is, certain Scott families. Scott families come from Scott's Isomorphism Theorem, which says that for a countable structure  $\mathcal{A}$ , there is an  $L_{\omega_1\omega}$  sentence whose countable models are exactly the isomorphic copies of  $\mathcal{A}$ . A *Scott family* for a structure  $\mathcal{A}$  is a countable family  $\Phi$  of  $L_{\omega_1\omega}$  formulas, possibly with finitely many fixed parameters from  $\mathcal{A}$ , such that:

- (i) Each finite tuple in  $\mathcal{A}$  satisfies some  $\psi \in \Phi$ ;

(ii) If  $\vec{a}, \vec{b}$  are tuples in  $\mathcal{A}$ , of the same length, satisfying the *same* formula in  $\Phi$ , then there is an automorphism of  $\mathcal{A}$  that maps  $\vec{a}$  to  $\vec{b}$ .

A *formally c.e. Scott family* is a c.e. Scott family consisting of finitary existential formulas. A *formally  $\Sigma_\alpha^0$  Scott family* is a  $\Sigma_\alpha^0$  Scott family consisting of computable  $\Sigma_\alpha$  formulas. Roughly speaking, computable infinitary formulas are  $L_{\omega_1\omega}$  formulas in which the infinite disjunctions and conjunctions are taken over computably enumerable (c.e.) sets. We can classify computable formulas according to their complexity in the following way: A computable  $\Sigma_0$  or  $\Pi_0$  formula is a finitary quantifier-free formula. Let  $\alpha > 0$  be a computable ordinal. A computable  $\Sigma_\alpha$  formula is a c.e. disjunction of formulas  $(\exists \vec{u})\theta(\vec{x}, \vec{u})$ , where  $\theta$  is computable  $\Pi_\beta$  for some  $\beta < \alpha$ . A computable  $\Pi_\alpha$  formula is a c.e. conjunction of formulas  $(\forall \vec{u})\theta(\vec{x}, \vec{u})$ , where  $\theta$  is computable  $\Sigma_\beta$  for some  $\beta < \alpha$ . Precise definition of computable infinitary formulas involves assigning indices to the formulas, based on Kleene's system of ordinal notations (see [2]). The important property of these formulas is given in the following theorem due to Ash.

**Theorem 1.1 (Ash)** *For a structure  $\mathcal{A}$ , if  $\theta(\vec{x})$  is a computable  $\Sigma_\alpha$  formula, then the set  $\{\vec{a} : \mathcal{A} \models \theta(\vec{a})\}$  is  $\Sigma_\alpha^0$  relative to  $\mathcal{A}$ .*

An analogous result holds for computable  $\Pi_\alpha$  formulas.

It is easy to see that if  $\mathcal{A}$  has a formally c.e. Scott family, then  $\mathcal{A}$  is relatively computably categorical. In general, if  $\mathcal{A}$  has a formally  $\Sigma_\alpha^0$  Scott family, then  $\mathcal{A}$  is relatively  $\Delta_\alpha^0$  categorical. Goncharov [10] showed that if  $\mathcal{A}$  is 2-decidable and computably categorical, then it has a formally c.e. Scott family. Ash [1] showed that, under certain decidability conditions on  $\mathcal{A}$ , if  $\mathcal{A}$  is  $\Delta_\alpha^0$  categorical, then it has a formally  $\Sigma_\alpha^0$  Scott family. For the relative notions, the decidability conditions are not needed. Ash, Knight, Manasse and Slaman [3] and Chisholm [6] independently proved that a computable structure  $\mathcal{A}$  is relatively  $\Delta_\alpha^0$  categorical iff it has a formally  $\Sigma_\alpha^0$  Scott family.

Cholak, Goncharov, Khoussainov and Shore [7] gave an example of a computable structure that is computably categorical, but ceases to be after naming any element of the structure. Such a structure is not relatively computably categorical. On the other hand, Millar [25] established that if a structure  $\mathcal{A}$  is 1-decidable, then any expansion of  $\mathcal{A}$  by finitely many constants remains computably categorical. Khoussainov and Shore [18] proved that there is a computably categorical structure without a formally c.e. Scott family whose expansion by any finite number of constants is computably categorical. A similar result was established by Kudinov by a quite different method. Using a modified family of enumerations constructed by Selivanov [29], Kudinov produced a computably categorical, 1-decidable structure without a formally c.e. Scott family.

A structure is *rigid* if it does not have nontrivial automorphisms. A computable structure is  $\Delta_\alpha^0$  *stable* if every isomorphism from  $\mathcal{A}$  onto a computable structure is  $\Delta_\alpha^0$ . If a computable structure is rigid and  $\Delta_\alpha^0$  categorical, then it

is  $\Delta_\alpha^0$  stable. A *defining family* for a structure  $\mathcal{A}$  is a set  $\Phi$  of formulas with one free variable and a fixed finite tuple of parameters such that:

- (i) Every element of  $A$  satisfies some formula  $\psi(x) \in \Phi$ ;
- (ii) No formula of  $\Phi$  is satisfied by more than one element of  $A$ .

A defining family  $\Phi$  is *formally*  $\Sigma_\alpha^0$  if it is a  $\Sigma_\alpha^0$  set of computable  $\Sigma_\alpha$  formulas. In particular, a defining family  $\Phi$  is *formally c.e.* if it is a c.e. set of finitary existential formulas. For a rigid computable structure  $\mathcal{A}$ , there is a formally  $\Sigma_\alpha^0$  Scott family iff there is a formally  $\Sigma_\alpha^0$  defining family.

It is not known whether for a computable limit ordinal  $\alpha$ , there is a computable structure that is  $\Delta_\alpha^0$  categorical but not relatively  $\Delta_\alpha^0$  categorical (see [13]). It is also not known whether for any computable successor ordinal  $\alpha$ , there is a rigid computable structure that is  $\Delta_\alpha^0$  categorical but not relatively  $\Delta_\alpha^0$  categorical. Another open question is whether every  $\Delta_1^1$  categorical computable structure must be relatively  $\Delta_1^1$  categorical (see [14]).

In Section 2, we investigate algorithmic properties of computable equivalence structures, their equivalence classes, and their characters. In Section 3, we examine effective categoricity of equivalence structures. We characterize the computably categorical equivalence structures and show that they are all relatively computably categorical. That is,  $\mathcal{A}$  is computably categorical if and only if the following two conditions are satisfied:

- (i) There is an upper bound on the size of the finite equivalence classes of  $\mathcal{A}$ ;
- (ii) There is at most one cardinal  $k$  such that  $\mathcal{A}$  has infinitely many equivalence classes of size  $k$ .

In Section 3, we characterize the relatively  $\Delta_2^0$  categorical equivalence structures as those with either finitely many infinite equivalence classes or with an upper bound on the size of the finite equivalence classes. We also consider the complexity of isomorphisms for structures  $\mathcal{A}$  and  $\mathcal{B}$  such that both  $\text{Fin}^\mathcal{A}$  and  $\text{Fin}^\mathcal{B}$  are computable or  $\Delta_2^0$ . Finally, we show that every computable equivalence structure is relatively  $\Delta_3^0$  categorical.

## 2 Computable Equivalence Structures

An equivalence structure  $\mathcal{A} = (A, E^A)$  consists of a set with a binary relation that is reflexive, symmetric and transitive. An equivalence structure  $\mathcal{A}$  is *computable* if  $A$  is a computable subset of  $\omega$  and  $E$  is a computable relation. If  $A$  is an infinite set (which is usual), we can assume, without loss of generality, that  $A = \omega$ . The  $\mathcal{A}$ -equivalence class of  $a \in A$  is

$$[a]^\mathcal{A} = \{x \in A : xE^\mathcal{A}a\}.$$

We generally omit the superscript  $\mathcal{A}$  when it can be inferred from the context.

We will proceed from the simpler structures, which are computably categorical, to the more complicated structures, which are  $\Delta_3^0$  categorical but not  $\Delta_2^0$  categorical.

**Definition 2.1** The character  $\chi(\mathcal{A})$  of an equivalence relation  $\mathcal{A}$  is the set  $\{\langle k, n \rangle : \mathcal{A} \text{ has at least } n \text{ equivalence classes of size } k\}$ .

We say that  $\mathcal{A}$  has *bounded character* if there is some finite  $K$  such that all finite  $\mathcal{A}$ -equivalence classes have size at most  $K$ .

The following lemmas will be needed.

**Lemma 2.2** For any computable equivalence structure  $\mathcal{A}$ :

- (a)  $\{\langle k, a \rangle : \text{card}([a]^{\mathcal{A}}) \leq k\}$  is a  $\Pi_1^0$  set and  $\{\langle k, a \rangle : \text{card}([a]^{\mathcal{A}}) \geq k\}$  is a  $\Sigma_1^0$  set.
- (b)  $\text{Inf}^{\mathcal{A}} = \{a : [a]^{\mathcal{A}} \text{ is infinite}\}$  is a  $\Pi_2^0$  set and  $\text{Fin}^{\mathcal{A}} = \{a : [a]^{\mathcal{A}} \text{ is finite}\}$  is a  $\Sigma_2^0$  set.
- (c)  $\chi(\mathcal{A})$  is a  $\Sigma_2^0$  set.

**Proof:** (a) The condition  $\text{card}([a]^{\mathcal{A}}) \leq k$  holds if and only if the statement

$$(\forall x_1) \cdots (\forall x_{k+1}) ( (x_1 E a \ \& \ \cdots \ \& \ x_{k+1} E a) \Rightarrow \bigvee_{i,j \neq k+1} x_i = x_j )$$

is satisfied.

(b) We have  $a \in \text{Fin}^{\mathcal{A}}$  if and only if

$$(\exists k)[\text{card}([a]^{\mathcal{A}}) = k]$$

and  $a \in \text{Inf}^{\mathcal{A}}$  if and only if  $a \notin \text{Fin}^{\mathcal{A}}$ .

(c) We have  $\langle k, n \rangle \in \chi(\mathcal{A})$  if and only if

$$(\exists x_1) \cdots (\exists x_n) ( \bigwedge_i \text{card}([x_i]) = k \ \& \ \bigwedge_{i \neq j} \neg(x_i E x_j) ).$$

□

We will say that a subset  $K$  of  $\omega \times \omega$  is a *character* if there is some equivalence structure with character  $K$ . This is the same as saying that for all  $n$  and  $k$ , if

$$\langle k, n+1 \rangle \in K \Rightarrow \langle k, n \rangle \in K.$$

**Lemma 2.3** For any  $\Sigma_2^0$  character  $K$ , there is a computable equivalence structure  $\mathcal{A}$  with character  $K$ , which has infinitely many infinite equivalence classes. Furthermore, in this structure,  $\{a : [a] \text{ is finite}\}$  is a  $\Pi_1^0$  set.

**Proof:** Let  $R$  be a computable relation such that

$$\langle k, n \rangle \in K \iff (\exists w)(\forall z)R(k, n, w, z).$$

We will define a set  $B$  of quadruples  $\langle k, n, w, z \rangle$  such that when we look at numbers only below  $z$ , we believe that  $w$  is the least witness that  $k, n \in K$ , and

such that for other initial segments below  $z$ , there is some  $v < w$  that could be such a witness. Define the set  $B$  as follows:

$$\begin{aligned} B = & \{ \langle k, n, w, z \rangle : (\forall y < z)(R(k, n, w, y)) \ \& \\ & (\forall v < w)(\exists y < z)(\neg R(k, n, v, y)) \& \\ & (\forall y < z)(\exists v < w)R(k, n, v, y) \}. \end{aligned}$$

The set  $B$  is a computable subset of  $\omega$  with an infinite complement. The equivalence structure  $\mathcal{A}$  will consist of one class for each element of  $B$ , together with an infinite family of infinite equivalence classes. Partition  $\omega \setminus B$  into two computable, infinite, disjoint subsets. Use the first subset to define the infinitely many infinite classes, and let the second subset be  $C = \{c_0, c_1, \dots\}$ . The classes with representatives from  $B$  are defined in stages. Let  $B = \{b_0, b_1, \dots\}$  and let  $b_i = \langle k_i, n_i, w_i, z_i \rangle$ .

At *stage* 0, we put  $\{c_0, \dots, c_{k_0-2}\}$  into the equivalence class of  $b_0$ .

After stage  $s$ , we have  $s$  equivalence classes with representatives  $b_0, \dots, b_{s-1}$ . For some  $i < s$ , the classes with representatives  $b_i$  have size  $k_i$ , and others have been declared to be infinite and have at least  $s$  elements. These partial classes contain elements  $c_0, \dots, c_{p(s)}$  from  $C$ .

At *stage*  $s+1$ , we check for all  $i \leq s$  and all  $z \leq s$  whether  $R(k_i, n_i, w_i, z)$ . If the class  $[b_i]$  has previously been declared to be infinite, we simply add one new element to this class. For any other  $i$  such that some  $R(k_i, n_i, w_i, z)$  fails with  $z < s$ , we declare  $[b_i]$  to be infinite and add  $s$  new elements from  $C$  to  $[b_i]$ . If  $b_{s+1}$  is not declared to be infinite, then we put  $k_{s+1} - 1$  elements from  $C$  into this class.

Let  $\mathcal{A}$  be the structure constructed by this process. It is clear that whenever  $\langle k, n \rangle \in K$ , then there will be a unique  $b_i = \langle k, n, w_i, z_i \rangle$  such that  $[b_i]$  has size  $k$  and that these are the only finite classes of  $\mathcal{A}$ . Thus,  $\chi(\mathcal{A}) = K$ . If  $\langle k, n \rangle \notin K$ , then the class  $[b_i]$  corresponding to  $\langle k, n \rangle$  will be infinite and we already have infinitely many infinite equivalence classes, so, clearly,  $\mathcal{A}$  has infinitely many infinite equivalence classes.

We note that for any pair  $(k, n)$ , there is at most one  $w$  such that for all  $z$ , we have  $(k, n, w, z) \in R$  and such that for some  $\tilde{z}$ , we have  $(k, n, w, \tilde{z}) \in B$ . Also, for each triple  $(k, n, w)$ , there is at most one  $z$  such that  $(k, n, w, z) \in B$ , and for each triple  $(k, n, z)$  there is at most one  $w$  such that  $(k, n, w, z) \in B$ . Now it is clear that  $[b_i]$  is finite if and only if  $(\forall z)R(k_i, n_i, w_i, z)$ , and  $[b_i]$  has  $k_i$  elements. Further, if  $k, n \in K$ , then

$$|\{[b_i] : |[b_i]| = k\}| = n.$$

Then for  $c \notin B$ ,  $[c]$  is finite if and only if

$$(\forall i)(cEb_i \Rightarrow [b_i] \text{ is finite}).$$

This completes the proof.  $\square$

**Lemma 2.4** *For any  $r \leq \omega$  and any bounded character  $K$  (whether  $K \in \Sigma_2^0$  or not), there is a computable equivalence structure  $\mathcal{A}$  with character  $K$ , which has exactly  $r$  infinite equivalence classes. Furthermore,  $\{a : [a]^{\mathcal{A}} \text{ is finite}\}$  is a computable set.*

**Proof:** The desired structure  $\mathcal{A}$  will have three components, each itself either finite or computable. First, there will be  $r$  infinite equivalence classes. Second, there will be a finite set  $\{k_1, \dots, k_m\} \subset \omega$  and an infinite family of equivalence classes of size  $k_i$  for  $i = 1, \dots, m$ . Third, there will be a finite set  $\{j_1, \dots, j_p\} \subset \omega$  with corresponding natural numbers  $n_i > 0$  for  $i = 1, \dots, p$  and  $j_i$  equivalence classes of size  $n_i$ . It is clear that a computable structure  $\mathcal{A}$  can be constructed such that each desired component is itself computable.  $\square$

The proof of Lemma 2.3 really needed the assumption of infinitely many infinite equivalence classes, since it is possible that either finitely many or infinitely many infinite equivalence classes come from  $B$ .

If there are just finitely many infinite equivalence classes, then the notion of  $s$ -functions and  $s_1$ -functions is important. These functions were introduced by Khisamiev in [16].

**Definition 2.5** *The function  $f : \omega^2 \rightarrow \omega$  is an  $s$ -function if the following hold:*

(i) *For every  $i$  and  $s$ ,  $f(i, s) \leq f(i, s + 1)$ .*

(ii) *For every  $i$ , the limit  $\lim_s f(i, s)$  exists.*

*Let  $m_i =_{\text{def}} \lim_s f(i, s)$ .*

*We say that  $f$  is an  $s_1$ -function if, in addition,  $m_i < m_{i+1}$  for each  $i$ .*

**Lemma 2.6** *Let  $\mathcal{A}$  be a computable equivalence structure with finitely many infinite equivalence classes and an infinite character. Then*

(i) *There exists a computable  $s$ -function  $f$  with corresponding limits  $m_i = \lim_s f(i, s)$  such that  $\langle k, n \rangle \in \chi(\mathcal{A})$  if and only if*

$$\text{card}(\{i : k = m_i\}) \geq n.$$

(ii) *If the character is unbounded, then there is a computable  $s_1$ -function  $g$  such that  $\mathcal{A}$  contains an equivalence class of size  $m_i$  for all  $i$ , where*

$$m_i = \lim_s g(i, s).$$

**Proof:** We may assume, without loss of generality, that  $\mathcal{A}$  has no infinite equivalence classes, since the infinite classes can be captured by a finite set of representatives.

(i) Define a computable sequence of representatives for all equivalence classes of  $\mathcal{A}$  by setting  $a_0 = 0$  and setting  $a_{i+1}$  to be the least  $a > a_i$  such that  $\neg(aEa_j)$  for all  $j \leq i$ . Now simply let

$$f(i, s) = \text{card}(\{a \leq s : aEa_i\}).$$

(ii) We will define a uniformly computable family  $a_i^s$  for  $i \leq s$  in such a way that  $a_i = \lim_s a_i^s$  converges. We will also define a computable sequence  $p_s$ , and let

$$f(i, s) = \text{card}(\{a \leq p_s : aEa_i^s\}).$$

Hence, we will have

$$m_i = \lim_s (\text{card}(\{a \leq s : aEa_i\}) = \text{card}([a_i])).$$

At stage 0, we have  $p_0 = 0$  and  $a_0^0 = 0$ , so that  $f(0, 0) = 1$ .

After stage  $s$ , we have  $p_s$  and  $a_0^s, \dots, a_s^s$  such that

$$f(i, s) = \text{card}(\{a \leq p_s : aEa_i^s\}),$$

and

$$f(0, s) < f(1, s) < \dots < f(s, s).$$

At stage  $s + 1$ , we look for the least  $p$  such that there is a sequence  $b_0, \dots, b_{s+1}$  with the property that

$$k_i = \text{card}(\{a \leq p : aEb_i\})$$

and  $k_0 < k_1 < \dots < k_{s+1}$ , and with the further property that  $b_i = a_i^s$  whenever there is no  $j \leq i$  and no  $aEa_j^s$  with  $p_s < a \leq p$ . Then we let  $a_i^{s+1} = b_i$  and  $p_{s+1} = p$ . To see that such  $p$  exists, simply let  $m$  be the largest such that  $[a_j^s] = \{a \leq p : aEa_j^s\}$  for all  $j \leq m$ , and let  $b_i = a_i^s$  for all  $i \leq m$ . Then use the fact that  $\chi(\mathcal{A})$  is unbounded to find  $b_{m+1}, \dots, b_{s+1}$  with

$$\begin{aligned} \text{card}([a_m^s]) &< \text{card}([b_{m+1}]) < \text{card}([b_{m+2}]) < \dots \\ &< \text{card}([b_{s+1}]), \end{aligned}$$

and take  $p$  large enough so that  $[b_i] = \{a \leq p : aEb_i\}$ . □

**Lemma 2.7** *For any computable  $s_1$ -function  $f$ , the range of  $f$  is a  $\Delta_2^0$  set.*

**Proof:** Let  $m_i = \lim_s f(i, s)$ . Since  $m_0 < m_1 < \dots$ , it follows that  $m \in \text{ran}(f)$  if and only if there exists  $i < m$  such that  $m = f(i)$ , which has the following two characterizations:

$$(\exists s)(\forall t > s)f(i, t) = m,$$

and

$$(\forall s)(\exists t > s)f(i, t) = m.$$

Thus, the range of  $f$  is both  $\Sigma_2^0$  and  $\Pi_2^0$ . □

**Lemma 2.8** *Let  $K$  be a  $\Sigma_2^0$  characteristic and let  $r$  be finite.*



- (i) Let  $f$  be a computable  $s$ -function with the corresponding limits  $m_i = \lim_s f(i, s)$  such that  $\langle n, k \rangle \in K$  if and only if

$$\text{card}(\{i : k = m_i\}) \geq n.$$

Then there is a computable equivalence structure  $\mathcal{A}$  with  $\chi(\mathcal{A}) = K$  and with exactly  $r$  infinite equivalence classes.

- (ii) Let  $f$  be a computable  $s_1$ -function with corresponding limits  $m_i = \lim_s f(i, s)$  such that  $\langle m_i, 1 \rangle \in K$  for all  $i$ . Then there is a computable equivalence structure  $\mathcal{A}$  with  $\chi(\mathcal{A}) = K$  and exactly  $r$  infinite equivalence classes.

**Proof:** Clearly, it suffices to prove the statements for  $r = 0$ . We may assume that  $f(i, 0) \geq 1$  for all  $i$ .

(i) Let  $a_i = 2i$ . We will build an equivalence structure  $\mathcal{A}$  with equivalence classes  $[a_i]$  of  $m_i$ . At stage 0, make the elements  $1, 3, \dots, 2(f(0, 0) - 1)$  equivalent to  $a_0$ . After stage  $s$ , we have exactly  $f(m, s)$  elements equivalent to  $a_m$  for each  $m \leq s$ . Then we add  $f(m, s + 1) - f(m, s)$  elements to  $[a_m]$  for  $m \leq s$  and put  $f(s + 1, s + 1) - 1$  elements into the class of  $a_{s+1}$ .

(ii) Since there is an  $s_1$ -function, the character  $K$  must be unbounded. We modify the argument for Lemma 2.3 as follows. The pool of elements to put into the equivalence classes is now simply  $\omega \setminus B$ .

Here is the first modification. When we find  $\neg R(k_i, n_i, w_i, z)$  for some  $z$  at stage  $s + 1$ , we can no longer create an infinite equivalence class, but we have already put  $k_i$  elements in the equivalence class of  $b_i$ . So we will set this bloc  $[b_i]$  aside until we find a number  $j$  and a stage  $s$  such that  $k_i \leq f(j, s)$ . Since there is an increasing sequence  $m_0 < m_1 < \dots$  corresponding to the  $s_1$ -function, such  $j$  and  $s$  will eventually be found. Then we will assign a marker  $j$  to  $b_i$  and add  $f(j, s) - k_i$  elements to the bloc to create an equivalence class with  $f(j, s)$  elements. If at a later stage  $t$  we have  $f(j, t) > f(j, s)$ , then we will add  $f(j, t) - f(j, s)$  more elements to the class. Since  $\lim_s f(j, s) = m_j$  converges, we will eventually have an equivalence class of size  $m_j$ .

This means that we may have created an extra equivalence class with  $f(j, s)$  elements, so the second modification is that when we create a class with  $k = f(j, s)$  elements, we may need (perhaps temporarily) to remove from our construction any class corresponding to  $\langle k, 1, w, z \rangle$ . That is, we set these (finitely many) blocs aside to be put into a larger class, just as if we had found that  $\neg R(k, n, w, z)$ , but we make a note that they may need to be revived later. If at some later stage  $t$ , we find  $k'$  such that

$$k' = f(j, t) > f(j, s) = k,$$

so that we will increase the size of the class with marker  $j$ , then we are going to remove the classes corresponding to  $\langle k', 1 \rangle$  and at the same time revive the classes corresponding to  $\langle k, 1 \rangle$ . At this stage, we remove the attachment to  $f(j, s)$  of the bloc and check to see for all  $b_i = \langle k, 1, w, z \rangle$ :

- (1) Whether  $R(k, 1, w, z)$  still holds for all  $z \leq t$ ;
- (2) Whether the bloc corresponding to  $b_i$  has been put into a larger class yet.

If (1) is false and (2) is true, then there is nothing else to do. If (1) and (2) are both false, then we keep the bloc aside for later use. If (1) is true and (2) is false, then we revive this bloc. If both (1) and (2) are true, then we create a new class with  $k$  elements for each  $\langle k, 1, w, z \rangle$  such that  $R(k, 1, w, z)$  still holds for all  $z \leq t$ .

We will now describe the construction in detail. Set  $C_{-1} = \omega - B$ . No  $j$ -markers are used at stage  $-1$ . If  $b_i = \langle k_i, n_i, w_i, z_i \rangle$  as in Lemma 2.3, let us say that  $b_i$  is *active* at stage  $s$  if for all  $z \leq s$ ,  $R(k_i, n_i, w_i, z)$ , and otherwise we say that  $b_i$  is *inactive*.

After  $s$  stages, we will have some equivalence classes with active representatives  $b_i = \langle k, n, w, z \rangle$  or revived representatives  $b'_i$ , containing  $k$  elements, such that for all  $z \leq s$ ,  $R(k, n, w, z)$ . We will have some equivalence classes containing  $f(j, s)$  elements corresponding to the  $s_1$ -function  $f$ . There will also be certain blocs of size  $k_i$  corresponding to inactive  $b_i$  and certain displaced blocs corresponding to active  $b_i$ , which are waiting to be put into a larger equivalence class. Finally, there are some active  $b_i = \langle k_1, 1, w, z \rangle$  that have been displaced by some equivalence class of size  $f(j, s) = k_i$  but may need to be revived. There is a pool  $C_s$  of remaining elements that may be used to fill out new equivalence classes. At stage  $s + 1$ , the following things may happen.

First, we check to see whether  $b_{s+1} = \langle k, n, w, z \rangle$  is active at stage  $s + 1$ . If so, then we check to see whether  $n = 1$  and  $k = f(j, s + 1)$  for some current equivalence class with marker  $j$ . If such  $j$  exists, then we put  $b_{s+1}$  into the pool  $C_{s+1}$ . Otherwise, we create an equivalence class with  $k$  elements consisting of  $b_{s+1}$  and  $k - 1$  elements from the pool  $C_s$ . If  $b_{s+1}$  is already inactive, then we simply add it to the pool  $C_{s+1}$ .

Second, we check for  $i \leq s$  whether some  $b_i$  that was active at stage  $s$  becomes inactive at stage  $s + 1$ . If such  $b_i$  was representing an equivalence class at stage  $s$ , then that class is set aside as a bloc to be attached to some  $f(j, t)$  at a later stage.

Third, we look for the smallest bloc that has been set aside for some inactive  $b_i$  at stage  $s$ , and check whether there exists a previously unused  $j \leq s + 1$  such that  $k_i \leq f(j, s + 1)$ . If so, then we create an equivalence class including this bloc and containing  $f(j, s + 1)$  elements.

Fourth, for all markers  $j$  that are being used at stage  $s$ , we check to see whether  $f(j, s + 1) > f(j, s)$ . If so, then we add  $f(j, s + 1) - f(j, s)$  elements to the corresponding equivalence class. We then displace any classes with representatives  $b_i$  for  $i \leq s$  such that  $k_i = f(j, s + 1)$  and  $n_i = 1$ . That is, we set aside this class as a bloc to be attached later to some  $f(j', t)$ . Finally, we revive any active  $b_i$  such that  $k_i = f(j, s)$  and  $n_i = 1$ , which were displaced by  $f(j, s)$ . This means that we create a completely new class with  $k_i$  elements and a new representative  $b'_i$  taken from the pool.

It is clear that eventually all elements from the pool are put into some equivalence class. It needs to be verified that this class eventually stabilizes at

some finite size  $k$  and that the resulting equivalence structure has the desired character  $\chi(\mathcal{A})$ .

Suppose that  $a$  is first put into some class attached to marker  $j$  at stage  $s$ . Then this class will have size  $f(j, t)$  at any later stage  $t$  and will stabilize with  $m_j$  elements. Next, suppose that  $a$  is first put into some bloc with representative  $b_i$  or  $b'_i$ . There are two cases. If  $b_i$  remains active at all stages and is never displaced by any  $f(j, t)$ , then this class has exactly  $k_i$  elements at all future stages. Otherwise, this class is set aside as a bloc at some later stage, and then eventually put into a class with some marker  $j$ , which will stabilize with  $m_j$  elements. This is guaranteed by the fact that there are infinitely many  $m_j > k_i$  and eventually the bloc containing  $b_i$  will have the highest priority. Thus, all equivalence classes stabilize at some finite size. Hence,  $\mathcal{A}$  has no infinite equivalence classes.

Now, fix a finite  $k$  and suppose that  $\langle k, n \rangle \in K$  for all  $n < r$ , where  $r \leq \infty$ . We need to verify that there are exactly  $r$  classes of size  $k$  in  $\mathcal{A}$ . For each  $n < r$ , there will be a unique representative  $b_i = \langle n, k, w, z \rangle$  that remains active at all stages, where  $w$  is the least such that  $(\forall z)R(n, k, w, z)$ . For  $n > 1$ , this  $b_i$  will represent a class of size  $k$  and can never be displaced. For  $n = 1$ , there are two possibilities. There can be some (unique) marker  $j$  such that  $m_j = k$ , which corresponds to a class stabilizing at size  $k$ . In this case, any classes corresponding to  $b_i$  (or any later  $b'_i$ ) will be displaced and eventually not revived (once  $f(j, s)$  converges to  $m_j$ ). On the other hand, if there is no such marker  $j$ , then eventually there will be a unique class with representative  $b_i$  (or some  $b'_i$ ) with  $k$  elements, which is never displaced. Classes represented by other  $b_p$  or by other markers can never have size  $k$ . Thus  $\chi(\mathcal{A}) = K$ .  $\square$

The necessity of the  $s_1$  function is seen by the following.

**Theorem 2.9** *There is an infinite  $\Delta_2^0$  set  $D$  such that for any computable equivalence structure  $\mathcal{A}$  with unbounded character  $K$  and no infinite equivalence classes,  $\{k : \langle k, 1 \rangle \in K\}$  is not a subset of  $D$ . Hence, for any  $s_1$  function  $f$  with  $m_0 < m_1 < \dots$ , where  $m_n = \lim_s f(n, s)$ , there exists  $i$  such that  $m_i \notin D$ .*

**Proof:** We use a method similar to Post's construction of a simple set. Let  $\mathcal{A}_e$  be the structure with universe  $\omega$  and relation  $E_e$  defined by

$$iE_e j \iff \langle i, j \rangle \in W_e$$

and let  $[a]_e = \{j : aE_e j\}$ . Then every computable equivalence structure is  $\mathcal{A}_e$  for some  $e$ , and  $[a]_e$  will be the equivalence class of  $a$ . Define a c.e. relation  $R$  by

$$R(e, a) \iff \text{card}([a]_e) > 2e.$$

Then by the standard uniformization theorem for c.e. relations (see Soare [31], p. 29), there exists a partial computable function  $f$ , called a *selector for  $R$* , such that, for every  $e$ ,

$$(\exists a)R(e, a) \Rightarrow R(e, f(e)).$$

Define  $D$  as follows.

$$k \in D \iff (\forall e < \frac{k}{2})(\text{card}([f(e)]_e) \neq k).$$

Then  $D$  is a  $\Delta_2^0$  set by part (a) of Lemma 2.2. For any  $\ell$ , the set

$$\hat{D} = \{n | (\exists x < \ell)(n = \text{card}([f(x)]_x))\}$$

has cardinality at most  $\ell$ , so that at most  $\ell$  of the elements from the set  $\{0, 1, \dots, 2\ell\}$  may be in  $\hat{D}$ . Thus, the complement of  $D$  contains at most  $e$  elements from  $\{0, 1, \dots, 2e\}$ . Hence, the complement of  $D$  is infinite. Now suppose that  $\mathcal{A}$  has unbounded character and has no infinite equivalence classes. Choose  $e$  so that  $\mathcal{A} = \mathcal{A}_e$ . Since  $\chi(\mathcal{A})$  is unbounded, there exists  $a$  such that  $R(e, a)$ , so that  $a = f(e)$ . Since  $\mathcal{A}$  has no infinite equivalence classes,

$$\text{card}([a]^{\mathcal{A}}) = \text{card}([f(e)]_e) = k > 2e$$

is finite. Then by definition,  $\langle k, 1 \rangle \in \chi(\mathcal{A})$  but  $k \notin D$ .

Now let  $f$  be any  $s_1$ -function, let  $m_i = \lim_s f(i, s)$ , and let  $K = \{\langle m_i, 1 \rangle : i \in \omega\}$ . Then there is an equivalence structure  $\mathcal{A}$  with character  $K$ . Therefore, some  $m_i \notin D$ .  $\square$

## 2.1 Computable categoricity of equivalence structures

We first investigate relative computable categoricity of computable equivalence structures by showing that they have a formally c.e. Scott family.

**Proposition 2.10** *If  $\mathcal{A}$  is a computable equivalence structure with only finitely many finite equivalence classes, then  $\mathcal{A}$  is relatively computably categorical.*

**Proof:** Choose parameters  $c_1, \dots, c_n$  which are representatives of the  $n$  finite equivalence classes. A Scott formula for any finite sequence  $\vec{a} = a_1, \dots, a_m$  of elements from  $A$  is a conjunction of two formulas. The first formula  $\phi(\vec{x})$  is the conjunction of all formulas  $x_i E x_j$  (when  $a_i E^{\mathcal{A}} a_j$ ) and  $\neg(x_i E x_j)$  (when it is not the case that  $a_i E^{\mathcal{A}} a_j$ ). The second formula  $\psi(\vec{x}, \vec{c})$  is the conjunction of all formulas  $x_i E c_j$  (when  $a_i E^{\mathcal{A}} c_j$ ) and  $\neg(x_i E c_j)$  (when it is not the case that  $a_i E^{\mathcal{A}} c_j$ ). It is clear that every tuple of elements from  $A$  satisfies one of these formulas.

Suppose that  $\vec{a}$  and  $\vec{b}$  satisfy the same Scott formula. Then, in particular, we have  $a_i E^{\mathcal{A}} a_j \iff b_i E^{\mathcal{A}} b_j$ .

For any tuple  $\vec{d}$ , the equivalence class  $[d_i]$  is finite if and only if some  $x_i E^{\mathcal{A}} c_j$  occurs in the Scott formula of  $\vec{d}$ , and  $[d_i]$  is infinite otherwise. We will define an automorphism  $H$  of  $\mathcal{A}$  mapping  $\vec{a}$  to  $\vec{b}$ .

For any equivalence class  $[a]$  containing none of the elements of  $\vec{a}$ , of  $\vec{b}$ , or of  $\vec{c}$ , the function  $H$  will simply be the identity map. We also define  $H(a_i) = b_i$  and  $H(c_i) = c_i$ . This induces a partial one-to-one map from the equivalence

classes of  $\mathcal{A}$  into the equivalence classes of  $\mathcal{A}$ , which fixes finite classes setwise and takes infinite classes to infinite classes. Within a particular finite class  $[a_i]$  of size  $n$ , the partial map from  $[a_i]$  to  $[b_i]$  defined on  $[a_i] \cap \{\vec{a}\}$  can be extended to an isomorphism of  $[a_i]$  onto  $[b_i]$ .

For the infinite classes (whether finitely or infinitely many) the partial isomorphism of the classes may similarly be extended to a total isomorphism of the classes. Likewise, the partial map taking  $a_i$  to  $b_i$  may be extended to map the infinite class  $[a_i]$  to the infinite class  $[b_i]$ .  $\square$

**Proposition 2.11** *Let  $\mathcal{A}$  be a computable equivalence structure with finitely many infinite classes, with bounded character and with at most one finite  $k$  such that there are infinitely many equivalence classes of size  $k$ . Then  $\mathcal{A}$  is relatively computably categorical.*

**Proof:** Let  $c_1, \dots, c_n$  be representatives for the finite classes not of size  $k$  and let  $d_1, \dots, d_p$  be representatives for the finitely many infinite classes. Then the Scott formula for a finite sequence  $\vec{a}$  from  $\mathcal{A}$  is the conjunction of three formulas, the first two as in the proof of Proposition 2.10, and the third is the conjunction of all formulas  $x_i E d_j$  (when  $a_i E^{\mathcal{A}} d_j$ ) and  $\neg(x_i E d_j)$  (when it is not the case that  $a_i E^{\mathcal{A}} d_j$ ). Then  $[a_i]$  is infinite if and only if  $a_i E^{\mathcal{A}} d_j$  for some  $j$ , and  $\text{card}([a_i]) = k$  if and only if  $\neg(a_i E^{\mathcal{A}} d_j)$  for all  $j$ , and also  $\neg(a_i E^{\mathcal{A}} c_j)$  for all  $j$ .

Suppose that  $\vec{a}$  and  $\vec{b}$  satisfy the same Scott formula. Then we can define an automorphism of  $\mathcal{A}$  extending the partial function which takes  $a_i$  to  $b_i$  as in the proof of Proposition 2.10.  $\square$

**Corollary 2.12** *Let  $\mathcal{A}$  be a computable equivalence structure of one of the following types:*

1.  $\mathcal{A}$  has only finitely many finite equivalence classes;
2.  $\mathcal{A}$  has finitely many infinite classes, has bounded character (i.e. only finitely many finite sizes of equivalence classes), and has at most one finite  $k$  such that there are infinitely many classes of size  $k$ .

*Then  $\mathcal{A}$  is relatively computably categorical.*

For structures  $\mathcal{A}$  with  $\text{Fin}^{\mathcal{A}}$  computable, there is a stronger result.

**Proposition 2.13** *Let  $\mathcal{A}$  and  $\mathcal{B}$  be isomorphic computable equivalence structures such that  $\text{Fin}^{\mathcal{A}}$  and  $\text{Fin}^{\mathcal{B}}$  are computable and such that  $\mathcal{A}$  has infinitely many equivalence classes of size  $k$  for at most one finite  $k$ . Then  $\mathcal{A}$  and  $\mathcal{B}$  are computably isomorphic.*

**Proof:** By Proposition 2.10,  $\text{Inf}^{\mathcal{A}}$  and  $\text{Inf}^{\mathcal{B}}$  are computably isomorphic and by Proposition 2.11,  $\text{Fin}^{\mathcal{A}}$  and  $\text{Fin}^{\mathcal{B}}$  are computably isomorphic.  $\square$

In the remainder of this section, we will show that no other equivalence structures are computably categorical. Here is the first case.

**Theorem 2.14** *Suppose that there exist  $k_1 < k_2 \leq \omega$  such that the computable equivalence structure  $\mathcal{A}$  has infinitely many equivalence classes of size  $k_1$  and infinitely many classes of size  $k_2$ . Then  $\mathcal{A}$  is not computably categorical.*

**Proof:** We will define structures  $\mathcal{C}$  and  $\mathcal{D}$  both isomorphic to  $\mathcal{A}$  and such that  $\{a : \text{card}([a]^{\mathcal{C}}) = k_1\}$  is a computable set, but

$$\{a : \text{card}([a]^{\mathcal{D}}) = k_1\}$$

is not computable. Then these two structures are not computably isomorphic, so  $\mathcal{A}$  is not computably categorical.

Observe that  $\chi(\mathcal{A})$  is a  $\Sigma_2^0$  set by part (c) of Lemma 2.2 and therefore

$$K = \chi(\mathcal{A}) \setminus \{\langle k_1, n \rangle : n < \omega\}$$

is also  $\Sigma_2^0$ . Thus, if  $\chi(\mathcal{A})$  is bounded, then there is a computable equivalence structure with character  $K$  and the same number of infinite equivalence classes as  $\mathcal{A}$ . If  $\chi(\mathcal{A})$  is unbounded, then by Lemma 2.6, there is an  $s_1$ -function  $f$  for  $\chi(\mathcal{A})$ . If  $k_1 \neq m_i$  for any  $i$ , then  $f$  will be an  $s_1$ -function for the character  $K$ . If  $k_1 = m_i$ , then define a new  $s_1$ -function  $g$  for  $K$  by  $g(j) = f(j)$  for  $i < j$ , and  $g(j) = f(j+1)$  for  $i \geq j$ . Then there is a computable structure  $\mathcal{B}$  with character  $K$  by Lemma 2.8. Now we can define a structure  $\mathcal{C} \simeq \mathcal{A}$  (that is, with character  $\chi(\mathcal{A})$ ) by setting

$$(2a+1) E^{\mathcal{C}} (2b+1) \iff a E^{\mathcal{B}} b,$$

and

$$(2(mk_1+i)) E^{\mathcal{C}} (2(nk_1+j)) \iff m = n,$$

where  $i, j < k_1$ . In this structure  $\mathcal{C}$ ,  $\{a : \text{card}([a]) = k_1\}$  is a computable set.

At the same time, we can build a structure  $\mathcal{D} \simeq \mathcal{A}$  such that  $\{a : \text{card}([a]^{\mathcal{D}}) = k_1\}$  is not computable. There are two cases, depending on whether  $k_2$  is finite.

First suppose that  $k_2$  is finite and build a computable structure  $\mathcal{C}'$  with character  $\{\langle k_1, n \rangle, \langle k_2, n \rangle : n < \omega\}$  in which  $\{a : \text{card}([a]) = k_1\}$  is a complete c.e. set, as follows. Let  $M$  be a complete c.e. set and note that  $M$  is both infinite and co-infinite. The equivalence classes of  $\mathcal{C}'$  will have representatives  $2i$  so that  $\text{card}([2i]) = k_1$  if  $i \notin M$ , and  $\text{card}([2i]) = k_2$  if  $i \in M$ . The odd numbers will act as a pool of elements to fill out the classes.

The construction of  $\mathcal{C}'$  is in stages. After stage  $s$ , there will be classes  $C_i^s$  containing  $2i$  which will have  $k_1$  elements if  $i \notin M_s$  and  $k_2$  elements if  $i \in M_s$ . At stage  $s+1$ , we add a new class containing  $2s+2$  and also containing  $k_1-1$  new odd elements from the pool if  $s+1 \notin M_{s+1}$  and containing  $k_2-1$  new elements from the pool if  $s+1 \in M_{s+1}$ . Also, for any  $i \leq s$  such that  $i \in M_{s+1} \setminus M_s$ , we will add  $k_2 - k_1$  new elements from the pool to the class with representative  $[2i]$ .

Next suppose that  $k_2 = \omega$ . Just modify the construction above so that when  $i \in M_s$ , the class  $[2i]$  contains  $\max\{k_1, s\}$  elements. The details are left to the reader.

Finally we combine  $\mathcal{A}$  and  $\mathcal{C}'$  into  $\mathcal{D}$  by coding  $\mathcal{A}$  on the odd numbers and  $\mathcal{C}'$  on the even numbers. Then

$$\text{card}([2a]) = k_1 \iff a \notin M,$$

so that  $\{d : \text{card}([d]) = k_1\}$  is not computable.  $\square$

We observe that in the proof of Theorem 2.14, if  $\text{Fin}^{\mathcal{A}}$  is computable, then for the case that  $k_2 < \omega$ ,  $\text{Fin}^{\mathcal{C}}$  and  $\text{Fin}^{\mathcal{D}}$  will also be computable. Thus we have the following.

**Proposition 2.15** *For any  $k_1 < k_2 < \omega$  and any computable equivalence structure  $\mathcal{A}$  with  $\text{Fin}^{\mathcal{A}}$  computable and with infinitely many equivalence classes of size  $k_1$  and infinitely many equivalence classes of size  $k_2$ , there is a computable equivalence structure  $\mathcal{B}$  isomorphic to  $\mathcal{A}$  with  $\text{Fin}^{\mathcal{B}}$  computable such that  $\mathcal{A}$  and  $\mathcal{B}$  are not computably isomorphic.*

For the next result, we want to consider the so-called isomorphism problem for a class of structures. For total recursive functions  $\phi_e : \omega \times \omega \rightarrow \{0, 1\}$ , let  $\mathcal{C}_e = (\omega, \equiv_e)$  be the structure with

$$m \equiv_e n \iff \phi_e(\langle m, n \rangle) = 1.$$

It is easy to check that  $\{e : \mathcal{C}_e \text{ is an equivalence structure}\}$  is a  $\Pi_2^0$  set. The following Lemma is immediate from Lemma 2.2.

**Convention:** We say that a set is  $D_3^0$  if it is a difference of two  $\Sigma_3^0$  sets.

**Lemma 2.16** (a) *For any finite  $r$ ,*

*$\{e : \mathcal{C}_e \text{ has at least } r \text{ infinite equivalence classes}\}$  is a  $\Sigma_3^0$  set.*

(b) *For any finite  $r$ ,  $\{e : \mathcal{C}_e \text{ has at exactly } r \text{ infinite equivalence classes}\}$  is a  $D_3^0$  set.*

(c)  *$\{e : \mathcal{C}_e \text{ has infinitely many infinite equivalence classes}\}$  is a  $\Pi_4^0$  set.*

We need to look at indices for  $\Sigma_2^0$  sets in general. Let  $\langle S_e : e \in \omega \rangle$  be an enumeration of the  $\Sigma_2^0$  sets, that is,

$$n \in S_e \iff (\exists m)(\langle m, n \rangle \notin W_e).$$

Then an enumeration  $\langle K_e : e \in \omega \rangle$  of the  $\Sigma_2^0$  characters may be defined by

$$\langle k, n \rangle \in K_e \iff (\forall j \leq n)(\langle k, j \rangle \in S_e).$$

**Lemma 2.17** *Let  $K$  be a fixed infinite  $\Sigma_2^0$  character. Then  $\{e : K_e = K\}$  is  $\Pi_3^0$  complete.*

**Proof:** The set  $\{e : K_e = K\}$  is certainly a  $\Pi_3^0$  set. Now, let  $P$  be an arbitrary  $\Pi_3^0$  set and let  $S$  be a  $\Sigma_2^0$  set such that

$$e \in P \iff (\forall k)(\langle k, e \rangle \in S),$$

where we may assume, without loss of generality, that

$$\langle k+1, e \rangle \in S \Rightarrow \langle k, e \rangle \in S.$$

Next, we define a computable function  $f$  such that

$$K_{f(e)} = K \iff e \in P.$$

Set

$$\langle k, n \rangle \in K_{f(e)} \iff \langle k, n \rangle \in K \ \& \ \langle k, e \rangle \in S \ \& \ \langle n, e \rangle \in S.$$

If  $e \in P$ , then  $\langle k, e \rangle \in S$  for all  $k$ , so that for every  $k$  and  $n$ ,

$$\langle k, n \rangle \in K_{f(e)} \iff \langle k, n \rangle \in K.$$

If  $e \notin P$ , then there is some  $k_0$  such that, for all  $k \geq k_0$ ,  $\neg S(k, e)$ . Since  $K$  is infinite, there is some  $\langle k, n \rangle \in K$  such that either  $k \geq k_0$  or  $n \geq k_0$ , and, therefore,  $\langle k, n \rangle \notin K_{f(e)}$ .  $\square$

We note that in the proof of Lemma 2.17,  $K_{f(e)} \subseteq K$  for all  $e$ .

**Theorem 2.18** *Let  $\mathcal{A}$  be a computable equivalence structure with character  $K$  such that there does not exist a computable equivalence structure  $\mathcal{B}$  with character  $K$  and with finitely many infinite equivalence classes. Then  $\{e : \mathcal{C}_e \simeq \mathcal{A}\}$  is  $\Pi_3^0$  complete.*

**Proof:** The set  $\{e : \mathcal{C}_e \simeq \mathcal{A}\}$  is  $\Pi_3^0$  by Lemmas 2.2 and 2.17, since  $\mathcal{C}_e \simeq \mathcal{A}$  if and only if  $\chi(\mathcal{C}_e) = K$ .

For the completeness, let the computable function  $f$  be as in the proof of Lemma 2.17. Use the technique of Lemma 2.3 uniformly to create the equivalence structure  $\mathcal{C}_{g(e)}$  with character  $K_{f(e)}$  and infinitely many infinite equivalence classes. Then  $\mathcal{C}_{g(e)}$  is isomorphic to  $K$  if and only if  $K_{f(e)} = K$ . The result now follows by Lemma 2.17.  $\square$

**Theorem 2.19** *Let  $K$  be an unbounded  $\Sigma_2^0$  character. Let  $\mathcal{A}$  be a computable equivalence structure with character  $K$  such that there does not exist a computable equivalence structure  $\mathcal{B}$  with character  $K$  and with finitely many infinite equivalence classes. Then  $\mathcal{A}$  is not computably categorical.*

**Proof:** If  $\mathcal{A} = (\omega, \equiv_{\mathcal{A}})$  is computably categorical, then  $\mathcal{C}_e \simeq \mathcal{A}$  if and only if  $\mathcal{A}$  and  $\mathcal{C}_e$  are computably isomorphic. But this has a  $\Sigma_3^0$  definition, that is,

$$(\exists a)[a \in Tot \ \& \ (\forall m)(\forall n)(m \equiv_e n \iff \phi_a(m) \equiv_{\mathcal{A}} \phi_a(n))].$$

This contradicts the  $\Pi_3^0$  completeness from Theorem 2.18.  $\square$

For characters with  $s_1$  functions, a structure may have finitely many or infinitely many infinite equivalence classes, and there is a higher complexity.



**Theorem 2.20** *Let  $\mathcal{A}$  be a computable equivalence structure with unbounded character  $K$  and with a finite number  $r$  of infinite equivalence classes.*

- (a) *If  $r = 0$ , then  $\{e : C_e \simeq \mathcal{A}\}$  is  $\Pi_3^0$  complete.*
- (b) *If  $r > 0$ , then  $\{e : C_e \simeq \mathcal{A}\}$  is  $D_3^0$  complete.*

**Proof:** (a) Suppose that  $\mathcal{A}$  has no infinite equivalence classes. Then  $\{e : C_e \simeq \mathcal{A}\}$  is a  $\Pi_3^0$  set, since  $C_e \simeq \mathcal{A}$  if and only if the following two facts hold:

- (1)  $\chi(C_e) = K$  (which is a  $\Pi_3^0$  condition by Lemma 2.17);
- (2)  $C_e$  has no infinite equivalence classes (which is a  $\Pi_3^0$  condition by Lemma 2.16).

For the completeness, let  $P$  be a given  $\Pi_3^0$  set. We construct a reduction of  $P$  to our set as follows. Let  $g$  be an  $s_1$ -function for  $K$ , let  $m_i = \lim_s g(i, s)$  and let  $f_0(i, s) = g(i, 2s)$  and  $f_1(i, s) = g(i, 2s + 1)$  so that  $f_0$  and  $f_1$  are both  $s_1$ -functions. Now let

$$K_1 = K \setminus \{\langle m_{2i}, n \rangle : i < \omega\}.$$

Let  $\phi$  be given by the proof of Lemma 2.17 so that  $K_{\phi(e)} = K_1$  if and only if  $e \in P$  and such that  $K_{\phi(e)} \subseteq K_1$  for all  $e$ . Then

$$K_{\phi(e)} \cup (K \cap \{\langle m_{2i}, n \rangle : i \in \omega\})$$

always has an  $s_1$ -function, so we can apply the proof of Lemma 2.8 to construct  $C_{\psi(e)}$  with character

$$K_{\phi(e)} \cup (K \cap \{\langle m_{2i}, n \rangle : i \in \omega\}),$$

which has no infinite equivalence classes. It is now clear that  $e \in P$  if and only if  $K_{\phi(e)} = K_1$ , which is if and only if  $C_{\psi(e)} \simeq \mathcal{A}$ .

Note that if we simply apply Lemma 2.17 to  $K$  itself, we find that  $K_{\phi(e)}$  is finite whenever  $K_{\phi(e)} \neq K$ , so that  $C_{\phi(e)}$  would also be finite, whereas we are assuming that all computable equivalence structures  $C_i$  have universe  $\omega$ .

(b) Suppose that  $\mathcal{A}$  has exactly  $r > 0$  infinite equivalence classes. Then  $\{e : C_e \simeq \mathcal{A}\}$  is a  $D_3^0$  set, that is, the difference of two  $\Sigma_3^0$  sets, since  $C_e \simeq \mathcal{A}$  if and only if both of the following two facts hold:

- (1)  $\chi(C_e) = K$  (which is a  $\Pi_3^0$  condition by Lemma 2.17);
- (2)  $C_e$  has exactly  $r$  infinite equivalence classes (which is a  $D_3^0$  condition by Lemma 2.16).

For the completeness, let  $P$  be a  $\Pi_3^0$  set as in part (a) and let  $Q$  be a  $\Sigma_3^0$  set. Now let  $R$  be a  $\Pi_2^0$  set such that, for all  $d$ ,

$$d \in Q \iff (\exists c)(\langle c, d \rangle \in R).$$

Without loss of generality, we may assume that when  $d \in Q$ , there exists a unique  $c$  such that  $\langle c, d \rangle \in R$ . It follows from the  $\Pi_2^0$  completeness of  $\{e : W_e \text{ is infinite}\}$  that there is a computable set  $T$  such that

$$\langle c, d \rangle \in R \iff (\{t : \langle c, d, t \rangle \in T\} \text{ is infinite}).$$

We will define a computable function  $\theta$  so that, for all  $d$  and  $e$ ,

$$C_{\theta(d,e)} \simeq \mathcal{A} \iff d \in Q \ \& \ e \in P.$$

$C_{\theta(d,e)}$  will be the disjoint union of three components.

The first component will be a structure  $\mathcal{B}$  that has no infinite equivalence classes and has character  $K_{\phi(e)}$ , where

$$e \in P \iff K_{\phi(e)} = K_1.$$

This is constructed as in part (a).

The second component,  $\mathcal{C}$ , is fixed for all  $e$ , has no infinite equivalence classes and has character

$$\{\langle m_{2i}, n \rangle : \langle m_{2i}, n+1 \rangle \in K\}.$$

This might be a finite structure.

The third component,  $\mathcal{D}$ , will have character  $\{\langle m_{2i}, 1 \rangle : i \in \omega\}$  and will have exactly  $r$  infinite equivalence classes if  $d \in Q$  and no infinite equivalence classes if  $d \notin Q$ . It suffices to give the argument when  $r = 1$  since then we can always take  $r$  copies of  $\mathcal{D}$  to get  $r$  infinite equivalence classes.

From the  $s_1$ -function  $f_0$  we create an infinite set of  $s_1$ -functions  $g_c$ , where

$$g_c(i, s) = f_0(2^c(2i+1)).$$

Let

$$m_{c,i} = \lim_s g_c(i, s).$$

Then  $\mathcal{D}$  will be the disjoint union of equivalence structures  $\mathcal{D}_c$  having character

$$K \cap \{\langle m_{c,i}, n \rangle : i, n \in \omega\},$$

and having exactly one infinite equivalence class if  $\langle c, d \rangle \in Q$  and no infinite equivalence class otherwise. It now suffices to construct  $\mathcal{D}_c$  with universe  $\omega$ .

Fix  $c$  and let  $n_i = m_{c,i}$ . The construction of the equivalence relation  $E$  on  $\mathcal{D}_c$  is in stages. At stage  $s$ , there will be equivalence classes  $C_i^s$  of size  $g_c(i, s)$  for all  $i < s$ . There will be a particular  $i = i^s$  such that

$$\{2t : \langle c, d, t \rangle \in T\} \subseteq C_{i^s}^s.$$

This class  $C_{i^s}^s$  is the *test* class. *Initially*, we have the empty structure. By stage  $s$ , all numbers  $< s$  will have been assigned to an equivalence class, and, hence, we will have decided whether  $aEb$  for all  $a, b < 2s$ .

At *stage*  $t+1$ , let  $i = i^t$  and check whether  $\langle c, d, t \rangle \in T$ . If it is, then we let  $i^{t+1} = t$  and we create the new class  $C_{i^{t+1}}^{t+1}$  by adding to  $C_{i^t}^t$  the element  $2t$ , along with  $g_c(t, t+1) - g_c(i, t) - 1$  new odd numbers. We also create a new class  $C_{i^t}^{t+1}$  with  $g_c(i, t+1) - g_c(i, t)$  new odd numbers. For all  $j$  such that  $j < t$  and  $j \neq i$ , we add  $g_c(j, t+1) - g_c(j, t)$  odd numbers to the class  $C_j^t$  to obtain  $C_j^{t+1}$ .

If  $\langle c, d, t \rangle \notin T$ , then for all  $j < t$ , we simply add  $g_c(j, t+1) - g_c(j, t)$  odd numbers to the class  $C_j^t$  to obtain  $C_j^{t+1}$  and we create the new class  $C^t$  with exactly  $g_c(t, t+1)$  new odd elements.

There are two possible outcomes of this construction. If  $\{t : \langle c, d, t \rangle \in T\}$  is finite, then after some stage  $t$ ,  $i^t$  becomes fixed and, thus, has limit  $i$ . Then for every  $i$ , the class  $C_i = \cup_t C_i^t$  will have exactly  $n_i$  elements, and every number will belong to one of these classes. Thus,  $\mathcal{D}_c$  has character  $\{\langle n_i, 1 \rangle : i \in \omega\}$  and has no infinite equivalence classes. If  $\{t : \langle c, d, t \rangle \in T\}$  is infinite, then  $\lim_t i^t = \infty$  and  $\mathcal{D}_c$  has one additional, infinite equivalence class, the test class, which is  $\cup_t C_{i^t}^t$ .

It follows that if  $d \notin Q$ , then each  $\mathcal{D}_c$  has character  $\{\langle m_{c,i}, 1 \rangle : i \in \omega\}$  and has no infinite equivalence classes, so that  $\mathcal{D}$  has character  $\{\langle m_{2i}, 1 \rangle : i \in \omega\}$  and has no infinite equivalence classes. If  $d \in Q$ , then one of the  $\mathcal{D}_c$  has one infinite equivalence class and the others have no infinite equivalence classes. Thus,  $\mathcal{D}$  has exactly one infinite equivalence class, as desired.  $\square$

**Theorem 2.21** *Let  $K$  be an unbounded  $\Sigma_2^0$  character, and let  $\mathcal{A}$  be a computable equivalence structure with character  $K$  and with finitely many infinite equivalence classes. Then  $\mathcal{A}$  is not computably categorical.*

**Proof:** This follows immediately from Theorem 2.20 as in the proof of Theorem 2.19  $\square$

Note that since there are finitely many infinite equivalence classes, for the structures  $\mathcal{A}$  and  $\mathcal{B}$  of Theorem 2.21 which are isomorphic but not computably isomorphic, both  $Fin^{\mathcal{A}}$  and  $Fin^{\mathcal{B}}$  are computable.

**Theorem 2.22** *Let  $\mathcal{A}$  be an equivalence structure with an unbounded character  $K$ , and with infinitely many infinite equivalence classes. Suppose that there exists an equivalence structure  $\mathcal{B}$  with character  $K$  and with finitely many infinite equivalence classes. Then  $\{e : \mathcal{C}_e \simeq \mathcal{A}\}$  is  $\Pi_4^0$  complete.*

**Proof:** It follows from Lemma 2.6 that  $K$  possesses an  $s_1$ -function  $g$ . Let  $m_i = \lim_s g(i, s)$ . Let  $f_0(i, s) = g(i, 2s)$  and  $f_1(i, s) = g(i, 2s + 1)$ , so that both  $f_0$  and  $f_1$  are  $s_1$ -functions. From the  $s_1$ -function  $f_0$  we create an infinite set of  $s_1$ -functions  $g_c(i, s)$ , where

$$g_c(i, s) = f_0(2^c(2i + 1)).$$

Set  $m_{c,i} = \lim_s g_c(i, s)$ . Let

$$K_0 = K \setminus \{\langle m_{2i}, n \rangle : i, n < \omega\},$$

and for each  $c$ , let

$$K_{c+1} = K \cap \{\langle m_{c,i}, n \rangle : i, n \in \omega\}.$$

Thus,  $K$  is the disjoint union of the characters  $K_c$ . Now,  $K_0$  has  $s_1$ -function  $f_1$ , and, therefore, there is a model  $\mathcal{A}_0$  with character  $K_0$  and no infinite equivalence classes.

Let  $P$  be a  $\Pi_4^0$  set. Let  $Q$  be a  $\Sigma_3^0$  relation such that

$$e \in P \iff (\forall c) Q(e, c).$$

We may assume that if  $e \notin P$ , then  $\{c : Q(e, c)\}$  is finite. Uniformizing the proof of Theorem 2.20, there is a computable binary function  $\phi$  such that  $\mathcal{C}_{\phi(e,c)}$  has character  $K_{c+1}$  for all  $e$  and  $c$ , and has exactly one infinite equivalence class if  $Q(e, c)$ , and no infinite equivalence class if  $\neg Q(e, c)$ .

Now define  $\mathcal{C}_{\psi(e)} = (\omega, E)$  as the effective union of the structures  $\mathcal{A}_0, \mathcal{C}_{\phi(e,c)}$ . That is, let  $E_0$  be the equivalence relation of  $\mathcal{A}_0$ , and let  $E_{e,c}$  be the equivalence relation of  $\mathcal{C}_{\phi(e,c)}$ . Let

$$E(2a, 2b) \iff E_0(a, b),$$

and for each  $c$ , let

$$E(2^c(2a+1), 2^c(2b+1)) \iff E_{e,c}(a, b);$$

for any other  $i, j$ , we let  $\neg E(i, j)$ . Then the structure  $\mathcal{C} = (\omega, E)$  clearly has character  $K = \cup_c K_c$ .

If  $e \in P$ , then  $Q(e, c)$  holds for all  $c$ , so each  $\mathcal{C}_{\phi(e,c)}$  has an infinite equivalence class. Hence,  $\mathcal{C}_{\psi(e)}$  has infinitely many infinite equivalence classes.

If  $e \notin P$ , then  $Q(e, c)$  holds for finitely many  $c$ , so finitely many  $\mathcal{C}_{\phi(e,c)}$  have exactly one infinite equivalence class, and the others have no infinite equivalence classes. Thus,  $\mathcal{C}_{\psi(e)}$  has finitely many infinite equivalence classes.

It follows that  $\mathcal{C}_{\psi(e)} \simeq \mathcal{A}$  if and only if  $e \in P$ .  $\square$

**Theorem 2.23** *Let  $\mathcal{A}$  be a computable equivalence structure with unbounded character  $K$  and with infinitely many infinite equivalence classes, such that there exists a computable equivalence structure  $\mathcal{B}$  with character  $K$  and with finitely many infinite equivalence classes. Then  $\mathcal{A}$  is not  $\Delta_2^0$  categorical.*

**Proof:** If  $\mathcal{A} = (\omega, \equiv_A)$  is  $\Delta_2^0$  categorical, then  $\mathcal{C}_e \simeq \mathcal{A}$  if and only if  $\mathcal{A}$  and  $\mathcal{C}_e$  are  $\Delta_2^0$  isomorphic. Thus, the set  $\{e : \mathcal{C}_e \simeq \mathcal{A}\}$  has a  $\Sigma_4^0$  definition. That is, with a c.e. complete set  $M$  as an oracle, we have

$$(\exists a) ([a \in Tot^M \& (\forall m)(\forall n)(m \equiv_e n \iff \phi_a^M(m) \equiv_A \phi_a^M(n))]).$$

This contradicts the  $\Pi_4^0$  completeness from Theorem 2.22.  $\square$

Combining these results, we obtain the following corollary.

**Corollary 2.24** *No equivalence structure with an unbounded character is computably categorical.*

We can now establish that for computable equivalence structures computable categoricity and relative computable categoricity coincide.

**Theorem 2.25** *All computably categorical equivalence structures are also relatively computably categorical.*

**Proof:** Suppose that  $\mathcal{A}$  is not relatively computably categorical and has character  $K$ . It follows from Propositions 2.10 and 2.11 that  $\mathcal{A}$  has infinitely many finite equivalence classes. First, suppose that  $K$  is bounded. Then there exists a finite  $k$  such that  $\mathcal{A}$  has infinitely many classes of size  $k$ . Now it follows from Proposition 2.11 that either  $\mathcal{A}$  has infinitely many infinite classes, or there are two finite numbers  $k_1$  and  $k_2$  such that  $\mathcal{A}$  has infinitely many classes of size  $k_1$  and infinitely many classes of size  $k_2$ . In either case, Theorem 2.14 implies that  $\mathcal{A}$  is not computably categorical.

Now, suppose that  $K$  is unbounded. Then there are two possibilities. Suppose first that  $K$  has no  $s_1$ -function. Then, by Theorem 2.19,  $\mathcal{A}$  is not computably categorical. Next, suppose that  $K$  has an  $s_1$ -function and that  $\mathcal{A}$  has infinitely many infinite equivalence classes. Then, by Theorem 2.23,  $\mathcal{A}$  is not computably categorical. Finally, suppose that  $\mathcal{A}$  has only finitely many infinite equivalence classes. Then  $\mathcal{A}$  is not computably categorical by Theorem 2.21.  $\square$

### 3 $\Delta_2^0$ Categoricity of equivalence structures

Next we continue with the analysis of  $\Delta_2^0$  categoricity.

**Theorem 3.1** *If  $\mathcal{A}$  is a computable equivalence structure with bounded character, then  $\mathcal{A}$  is relatively  $\Delta_2^0$  categorical.*

**Proof:** Let  $k$  be the maximum size of any finite equivalence class. The key fact here is that  $[a]$  is infinite if and only if  $[a]$  contains at least  $k + 1$  elements, which is a  $\Sigma_2^0$  condition. By Lemma 2.2, there is a  $\Delta_2^0$  formula which characterizes the elements  $a$  with a finite equivalence class of size  $m$ . Then a Scott formula for the tuple  $\langle a_1, \dots, a_m \rangle$  includes a formula  $\psi_i(x_i)$  for each  $a_i$ , giving the cardinality of  $[a_i]$ , together with formulas  $\psi_{i,j}(x_i, x_j)$  for each  $i$  and  $j$  which express whether  $a_i E a_j$ . It now follows, as in the proof of Proposition 2.10, that whenever  $\vec{a}$  and  $\vec{b}$  have the same Scott formula, then there is an automorphism of  $\mathcal{A}$  taking  $\vec{a}$  to  $\vec{b}$ . Thus,  $\mathcal{A}$  is relatively  $\Delta_2^0$  categorical.  $\square$

**Theorem 3.2** *If  $\mathcal{A}$  is a computable equivalence structure with finitely many infinite equivalence classes, then  $\mathcal{A}$  is relatively  $\Delta_2^0$  categorical.*

**Proof:** There is a  $\Sigma_1^0$  Scott formula for each element with an infinite equivalence class, by the proof of Proposition 2.11. There is a  $\Sigma_1^0$  Scott formula for each element with a finite equivalence class, by the proof of Theorem 3.1. It now follows, as before, that  $\mathcal{A}$  is relatively  $\Delta_2^0$  categorical.  $\square$

This leads to a stronger result for structures  $\mathcal{A}$  with  $Fin^{\mathcal{A}}$  computable.

**Theorem 3.3** *For any two isomorphic computable equivalence structures  $\mathcal{A}$  and  $\mathcal{B}$  such that  $Fin^{\mathcal{A}}$  and  $Fin^{\mathcal{B}}$  are both computable,  $\mathcal{A}$  and  $\mathcal{B}$  are  $\Delta_2^0$  categorical.*

**Proof:** It follows from Proposition 2.10 that  $Inf^{\mathcal{A}}$  and  $Inf^{\mathcal{B}}$  are computably isomorphic and it follows from Theorem 3.2 that  $Fin^{\mathcal{A}}$  and  $Fin^{\mathcal{B}}$  are  $\Delta_2^0$  isomorphic. Now the two mappings may be combined to define a  $\Delta_2^0$  isomorphism between  $\mathcal{A}$  and  $\mathcal{B}$  since  $Fin^{\mathcal{A}}$  and  $Fin^{\mathcal{B}}$  are computable.  $\square$

In fact, we observe that this result still holds if we only assume that  $Fin^{\mathcal{A}}$  and  $Fin^{\mathcal{B}}$  are both  $\Delta_2^0$ .  $Inf^{\mathcal{A}}$  and  $Inf^{\mathcal{B}}$  are still  $\Delta_2^0$  isomorphic and there is a  $\Delta_2^0$  enumeration of the finite equivalence classes of  $\mathcal{A}$  and  $\mathcal{B}$  which will induce a  $\Delta_2^0$  isomorphism between  $Fin^{\mathcal{A}}$  and  $Fin^{\mathcal{B}}$ .

**Theorem 3.4** *Let  $\mathcal{A}$  be a computable equivalence structure with infinitely many infinite equivalence classes, and with unbounded character which has a computable  $s_1$ -function. Then  $\mathcal{A}$  is not  $\Delta_2^0$ -categorical.*

**Proof:** We will build  $\mathcal{B}_1 \simeq \mathcal{B}_2 \simeq \mathcal{A}$  in such a way as to diagonalize against  $\Delta_2^0$  isomorphisms between  $\mathcal{B}_1$  and  $\mathcal{B}_2$ . Let  $\varphi_e^{\Delta_2^0}$  be a computable enumeration of all partial functions which are computable with a  $\Delta_2^0$  oracle. We will seek to meet the following requirements:

$$R_e: \quad \forall x \varphi_e^{\Delta_2^0} \downarrow \Rightarrow \exists x_e \left| \left[ \varphi_e^{\Delta_2^0}(x_e) \right]^{\mathcal{B}_2} \right| \neq \left| [x_e]^{\mathcal{B}_1} \right|$$

We will construct  $\mathcal{B}_i$  exactly as in Lemma 2.3, with the following exceptions. We begin by partitioning  $\omega$  into two infinite disjoint parts. One will provide ordinary elements for the domain. The other we will enumerate by  $\{x_e\}_{e \in \omega}$ , and so  $x_e$  will serve as a witness for  $R_e$ . Also, in place of the elements  $(k, n, w, z)$  elements, we will use elements of the form  $(k, n, w, z, q)$ , where all except the last coordinate work just as before. Until permission is given to use elements where  $q > \tilde{q}$ , no such elements will be used by the construction.

At stage  $s$ , we say that  $R_e$  requires attention if  $x_e \leq s$  and  $\forall x \leq s \varphi_e^{\Delta_2^0}(x) \downarrow \wedge \left| \left[ \varphi_e^{\Delta_2^0}(x_e) \right]^{\mathcal{B}_2} \right| \neq \left| [x_e]^{\mathcal{B}_1} \right|$ . For every  $e$  that requires attention at stage  $s$ , we will act in the following way. We will want to arrange that  $\left[ \varphi_e^{\Delta_2^0}(x_e) \right]^{\mathcal{B}_2}$  is of some finite size, and that  $[x_e]^{\mathcal{B}_1}$  is larger.

To achieve this, we will add elements to  $\left[ \varphi_e^{\Delta_2^0}(x_e) \right]^{\mathcal{B}_2}$ , following the  $s_1$ -function, as in Lemma 2.8. All other instructions about adding elements to  $\left[ \varphi_e^{\Delta_2^0}(x_e) \right]^{\mathcal{B}_2}$  will now be discarded. If  $\left[ \varphi_e^{\Delta_2^0}(x_e) \right]^{\mathcal{B}_2}$  contained an element of the form  $(k, n, w, z, q)$ , we will give permission to the element  $(k, n, w, z, q+1)$ , and will catch it up to the point reached by  $(k, n, w, z, q)$ .

In  $\mathcal{B}_1$ , we will choose a  $j$  larger than that used for the input of the  $s_1$ -function for  $\left[ \varphi_e^{\Delta_2^0}(x_e) \right]^{\mathcal{B}_2}$ . Increase the size of  $[x_e]^{\mathcal{B}_1}$ , following the  $s_1$ -function as in Lemma 2.8. If  $[x_e]^{\mathcal{B}_1}$  contained an element of the form  $(k, n, w, z, q)$ , we will give permission to the element  $(k, n, w, z, q+1)$ , and will catch it up to the point reached by  $(k, n, w, z, q)$ .

Now by the proofs of Lemmas 2.3 and 2.8, the structures  $\mathcal{B}_1$  and  $\mathcal{B}_2$  are equivalence structures with infinitely many infinite equivalence classes, and with

character  $\chi(\mathcal{A})$ . It remains to show that they are not  $\Delta_2^0$  isomorphic.

Suppose that  $\varphi_e^{\Delta_2^0}$  is an isomorphism from  $\mathcal{B}_1$  to  $\mathcal{B}_2$ . Suppose first that  $[x_e]^{\mathcal{B}_1}$  is finite, of size  $n$ . There was some stage  $s$  at which  $|[x_e]^{\mathcal{B}_1}| = n$ , and at which  $R_e$  received attention. This stage prevented  $\varphi_e^{\Delta_2^0}$  from being an isomorphism.

Now suppose that  $[x_e]^{\mathcal{B}_1}$  is infinite. Now  $R_e$  received attention infinitely often, which could only happen if  $\varphi_e^{\Delta_2^0}(x_e) \uparrow$ .  $\square$

It remains to consider the case of an unbounded character  $K$  with no  $s_1$ -function. Recall from Lemma 2.3 that we may construct a computable equivalence structure  $\mathcal{A}$  with  $Fin^{\mathcal{A}}$  a  $\Pi_1^0$  set. If we could also construct a computable equivalence structure  $\mathcal{B}$  with  $Fin^{\mathcal{B}}$  not a  $\Delta_2^0$  set, then it would follow that  $\mathcal{A}$  is not  $\Delta_2^0$  isomorphic to  $\mathcal{B}$ . Surprisingly, we cannot make  $Fin^{\mathcal{B}}$  a complete  $\Sigma_2^0$  set, as we could when  $K$  had an  $s_1$ -function. This is because of the following result.

**Theorem 3.5** *Let  $\mathcal{A}$  be a computable equivalence structure and let  $C$  be an infinite c.e. subset of  $Fin^{\mathcal{A}}$ . Then  $\mathcal{A}$  possesses a computable  $s$ -function  $f$ . Furthermore, if  $\{card([c]) : c \in C\}$  is unbounded, then  $\mathcal{A}$  possesses a computable  $s_1$ -function  $f$ . Thus, there is a computable structure with character  $\chi(\mathcal{A})$  and with no infinite equivalence classes.*

**Proof:** Let  $\mathcal{A} = (\omega, E)$  where  $E$  is a computable relation. Let

$$C_1 = \{a : (\exists c \in C) cEa\}.$$

Then  $C_1$  is also an infinite c.e. set. Fix its computable enumeration  $\{c_0, c_1, \dots\}$  of  $C_1$ , without repetition. Now let

$$f(i, s) = card(\{x \leq s : xEc_i\}).$$

Let  $\mathcal{A}_1 = (\omega, E_1)$  where  $iE_1j$  if and only if  $c_iEc_j$  and let  $K_1 = \chi(\mathcal{A}_1)$ . Then  $f$  is clearly an  $s$ -function for  $K_1$ , and  $K_1 \subset \chi(\mathcal{A})$ . It follows from Lemma 2.8 that there is a computable structure with character  $\chi(\mathcal{A})$  and with no infinite equivalence classes.

Now, suppose that  $\{card([c]) : c \in C\}$  is unbounded. Then  $\mathcal{A}_1$  has unbounded character and no infinite equivalence classes. Thus,  $K_1$  possesses an  $s_1$ -function  $g$  by Lemma 2.6 and hence  $K$  possesses the same  $s_1$ -function.  $\square$

**Theorem 3.6** *If the unbounded character  $K$  has no  $s_1$ -function, then there is no computable equivalence structure  $\mathcal{A}$  with character  $K$  such that  $Fin^{\mathcal{A}}$  is  $\Sigma_2^0$  complete, or even  $\Sigma_1^0$  hard.*

**Proof:** Let  $M$  be a complete c.e. set and suppose that there were a computable function  $f$  such that

$$i \in M \iff f(i) \in Fin^{\mathcal{A}}.$$

Then  $C = \{f(i) : i \in M\}$  is a c.e. subset of  $Fin^{\mathcal{A}}$ . If  $C$  is finite, say  $C = \{c_1, \dots, c_t\}$ , then

$$i \in M \iff (f(i) = c_1 \vee f(i) = c_2 \vee \dots \vee f(i) = c_t),$$

so that  $M$  would be a computable set. Thus,  $C$  is infinite. Now suppose that  $\{card([c]) : c \in C\}$  is bounded by some finite  $k$ . Then  $C$  is a subset of the  $\Pi_1^0$  set  $P$ , where

$$P = \{a : card([a]) \leq k\}.$$

Since we have

$$i \in M \iff f(i) \in P,$$

that would imply that  $M$  is a  $\Pi_1^0$  set. This contradiction shows that  $\{card([c]) : c \in C\}$  is unbounded. It now follows by Theorem 3.5 that  $K$  possesses an  $s_1$ -function.  $\square$

**Open Question:** Let  $\mathcal{A}$  be a computable equivalence structure having unbounded character, infinitely many infinite equivalence classes, and no  $s_1$ -function. Further assume that  $Fin^{\mathcal{A}}$  is Turing incomparable with  $\emptyset'$ . Does such a structure exist, and if so, is this structure  $\Delta_2^0$  categorical?

Theorem 3.6 may provide some evidence that such a structure, if it exists, may in fact be  $\Delta_2^0$  categorical. Nevertheless, we can show that such a structure cannot be *relatively*  $\Delta_2^0$  categorical.

**Theorem 3.7** *If the computable equivalence structure  $\mathcal{A}$  has unbounded character and infinitely many infinite equivalence classes, then it is not relatively  $\Delta_2^0$  categorical.*

**Proof:** Suppose, on the contrary, that an element  $a$  with an infinite equivalence class had a  $\Sigma_2^0$  Scott formula  $\psi(x, \vec{d})$ . Since there are only finitely many parameters  $\vec{d}$  involved, we may assume that  $[a]$  does not contain any of the parameters  $\vec{d}$ . (This is where we use the assumption that there are infinitely many infinite equivalence classes.) Then by choosing elements  $c_1, \dots, c_n$  of  $\mathcal{A}$  to instantiate the existentially quantified variables in  $\psi(x, \vec{d})$ , we would have a computable  $\Pi_1^0$  formula  $\theta(x, \vec{d}, \vec{c})$ , where  $\vec{c} = c_1, \dots, c_n$ , that is satisfied by  $a$ .

Now, it is easy to see that for *any* submodel  $\mathcal{M}$  of  $\mathcal{A}$  that contains  $a$ ,  $\vec{d}$  and  $\vec{c}$ , we have  $\mathcal{M} \models \theta(a, \vec{d}, \vec{c})$ . In fact, since our structures are relational,  $\mathcal{A} \models \theta(b, \vec{d}, \vec{c})$  iff and only if  $\mathcal{M} \models \theta(b, \vec{d}, \vec{c})$  for all finite submodels  $\mathcal{M}$  of  $\mathcal{A}$  that contain  $b$ ,  $\vec{d}$  and  $\vec{c}$ .

Thus, in particular, for the finite subset  $C = \{a, \vec{c}, \vec{d}\}$  of  $\omega$ , we have  $\mathcal{C} = (C, E^{\mathcal{C}}) \models \theta(a, \vec{d}, \vec{c})$ . Suppose that  $\vec{c}$  contains  $m \leq n$  elements of  $[a]$  and choose  $b$  such that  $[b] \cap C = \emptyset$  and  $m < card([b]) < \omega$ . (Here we use the fact that the character of  $\mathcal{A}$  is unbounded.) Let  $\mathcal{B} \simeq \mathcal{C}$  contain  $m$  elements of  $[b]$  (including  $b$ ), together with  $C \setminus [a]$ . Let  $\vec{e}$  denote the image of  $\vec{c}$  under the isomorphism between  $\mathcal{C}$  and  $\mathcal{B}$ . Then  $\mathcal{B} \models \theta(b, \vec{d}, \vec{e})$ . Furthermore, let  $\mathcal{B}'$  be any finite submodel of  $\mathcal{A}$  such that  $\mathcal{B} \subseteq \mathcal{B}'$ . Then it is easy to extend  $\mathcal{C}$  to a finite submodel  $\mathcal{C}'$  that is isomorphic to  $\mathcal{B}'$  (where the isomorphism fixes  $\vec{d}$



pointwise and takes  $a$  to  $b$ ). Thus,  $\mathcal{B}' \models \theta(b, \vec{d}, \vec{e})$  as well. It follows that  $\mathcal{A} \models \theta(b, \vec{d}, \vec{e})$ . Hence,  $\mathcal{A} \models \psi(b, \vec{d})$ .

But there certainly can be no automorphism of  $\mathcal{A}$  mapping  $a$  to  $b$ , since  $[a]$  is infinite and  $[b]$  is finite. Thus, in fact,  $a$  cannot have a  $\Sigma_2^0$  Scott formula.  $\square$

We conclude this section by looking at  $\Delta_3^0$  categoricity.

**Theorem 3.8** *Every computable equivalence structure is relatively  $\Delta_3^0$  categorical.*

**Proof:** Any element with an infinite equivalence class has a  $\Pi_2^0$  Scott formula, while the other elements even have  $\Delta_2^0$  Scott formulas. Thus, every tuple  $\langle a_1, \dots, a_n \rangle$  has a  $\Sigma_3^0$  Scott formula.  $\square$

## References

- [1] C.J. Ash, “Categoricity in hyperarithmetical degrees,” *Annals of Pure and Applied Logic* 34 (1987), pp. 1–14.
- [2] C.J. Ash and J. F. Knight, *Computable Structures and the Hyperarithmetical Hierarchy* (Elsevier, Amsterdam, 2000).
- [3] C. Ash, J. Knight, M. Manasse and T. Slaman, “Generic copies of countable structures,” *Annals of Pure and Applied Logic* 42 (1989), pp. 195–205.
- [4] C.J. Ash and A. Nerode, “Intrinsically recursive relations,” in *Aspects of Effective Algebra*, ed. by J. N. Crossley, Upside Down A Book Co., Steel’s Creek, Australia, 1981, pp. 26–41.
- [5] D. Cenzer and J. Remmel, “Polynomial-time Abelian groups,” *Annals of Pure and Applied Logic* 56 (1992), pp. 313–363.
- [6] J. Chisholm, “Effective model theory vs. recursive model theory,” *Journal Symbolic Logic* 55 (1990), pp. 1168–1191.
- [7] P. Cholak, S. Goncharov, B. Khoussainov and R.A. Shore, “Computably categorical structures and expansions by constants,” *Journal of Symbolic Logic* 64 (1999), pp. 13–37.
- [8] R.G. Downey, “Computability theory and linear orderings,” in: Yu. L. Ershov, S.S. Goncharov, A. Nerode and J.B. Remmel, editors, *Handbook of Recursive Mathematics*, vol. 2 (North-Holland, Amsterdam, 1998), pp. 823–976.
- [9] S.S. Goncharov, “Autostability of models and abelian groups,” *Algebra and Logic* 19 (1980), pp. 23–44 (Russian), pp. 13–27 (English translation).

- [10] S.S. Goncharov, “The quantity of non-autoequivalent constructivizations,” *Algebra and Logic* 16 (1977), pp. 257–282 (Russian), pp. 169–185 (English translation).
- [11] S.S. Goncharov, “Autostability and computable families of constructivizations,” *Algebra and Logic* 14 (1975), pp. 647–680 (Russian), pp. 392–409 (English translation).
- [12] S.S. Goncharov and V.D. Dzgoev, “Autostability of models,” *Algebra and Logic* 19 (1980), pp. 45–58 (Russian), pp. 28–37 (English translation).
- [13] S.S. Goncharov, V. S. Harizanov, J. F. Knight, C.F.D. McCoy, R. G. Miller, and R. Solomon, “Enumerations in computable structure theory,” to appear in the *Annals of Pure and Applied Logic*.
- [14] S.S. Goncharov, V.S. Harizanov, J.F. Knight and R.A. Shore, “ $\Pi_1^1$  relations and paths through  $\mathcal{O}$ ,” *Journal of Symbolic Logic* 69 (2004), pp. 585–611.
- [15] S. Goncharov, S. Lempp and R. Solomon, “The computable dimension of ordered abelian groups,” *Advances in Mathematics* 175 (2003), pp. 102–143.
- [16] N.G. Khisamiev, “Constructive Abelian  $p$ -groups ” *Siberian Advances in Mathematics* 2 (1992), pp. 68–113
- [17] N.G. Khisamiev, “Constructive Abelian groups,” in: Yu. L. Ershov, S.S. Goncharov, A. Nerode and J.B. Remmel, editors, *Handbook of Recursive Mathematics*, vol. 2 (North-Holland, Amsterdam, 1998), pp. 1177–1231.
- [18] B. Khoussainov and R.A. Shore, “Computable isomorphisms, degree spectra of relations and Scott families,” *Annals of Pure and Applied Logic* 93 (1998), pp. 153–193.
- [19] O.V. Kudinov, “An autostable 1-decidable model without a computable Scott family of  $\exists$ -formulas,” *Algebra and Logic* 35 (1996), pp. 458–467 (Russian), pp. 255–260 (English translation).
- [20] O.V. Kudinov, “A description of autostable models,” *Algebra and Logic* 36 (1997), pp. 26–36 (Russian), pp. 16–22 (English translation).
- [21] P. LaRoche, “Recursively presented Boolean algebras,” *Notices AMS* 24 (1977), A552–A553.
- [22] S. Lempp, C.F.D. McCoy, R.G. Miller and D. R. Solomon, “Computable categoricity of trees of finite height,” *Journal of Symbolic Logic* 70 (2005), pp. 151–215.
- [23] C.F.D. McCoy, “ $\Delta_2^0$ -categoricity in Boolean algebras and linear orderings,” *Annals of Pure and Applied Logic* 119 (2003), pp. 85–120.

- [24] G. Metakides and A. Nerode, “Effective content of field theory,” *Annals of Mathematical Logic* 17 (1979), pp. 289–320.
- [25] T. Millar, “Recursive categoricity and persistence,” *Journal of Symbolic Logic* 51 (1986), pp. 430–434.
- [26] R. Miller, “The computable dimension of trees of infinite height,” to appear in the *Journal of Symbolic Logic*.
- [27] A.T. Nurtazin, “Strong and weak constructivizations and computable families,” *Algebra and Logic* 13 (1974), pp. 311–323 (Russian), pp. 177–184 (English translation).
- [28] J.B. Remmel, “Recursively categorical linear orderings,” *Proceedings of the American Mathematical Society* 83 (1981), pp. 387–391.
- [29] V.L. Selivanov, “Numerations of families of general recursive functions,” *Algebra and Logic* 15 (1976), pp. 205–226 (Russian), pp. 128–141 (English translation).
- [30] R.L. Smith, “Two theorems on autostability in p-groups,” in *Logic Year 1979–80, Univ. Connecticut, Storrs*, Lecture Notes in Mathematics 859, Springer, Berlin (1981), pp. 302–311.
- [31] R.I. Soare, *Recursively Enumerable Sets and Degrees. A Study of Computable Functions and Computably Generated Sets* (Springer-Verlag, Berlin, 1987).